PURINA® ANIMAX®



Specie: Many Type of Feed: Supplement Form of Feed: Pellet

General Description: A nutritious, pelleted feed supplement formulated for many different livestock species.

PURINA® ANIMAX®

FOR HORSE, DAIRY CALVES AND MILK COWS, BEEF, TURKEY, SWINE, CHICKEN, PHEASANT, QUAIL, PARTRIDGE, DUCK, RABBIT & GOAT.

GUARANTEED ANALYSIS

Crude Protein, (Min)	28.00 %
Lysine, (Min)	1.50 %
Methionine, (Min)	0.40 %
Crude Fat, (Min)	
Crude Fiber, (Min)	3.00 %
Crude Fiber, (Max)	7.00 %
Acid Detergent Fiber (ADF), (Max)	
Neutral Detergent Fiber (NDF), (Max)	
Ash, (Max)	11.00 %
Calcium (Ca), (Min)	1.25 %
Calcium (Ca), (Max)	1.75 %
Phosphorus (P), (Min)	0.90 %
Salt (NaCl), (Min)	
Salt (NaCl), (Max)	
Sodium (Na), (Max)	
Potassium (K), (Min)	
Copper (Cu), ppm, (Min)	
Copper (Cu), ppm, (Max)	
Selenium (Se), ppm, (Min)	
Zinc (Zn), (Min)	140
Vitamin A, I.U./Ib, (Min)	30000
MODEDIENTO	

INGREDIENTS Note: ingredients differ by manufacturing plant

Dehulled Soybean Meal, Wheat Middlings, Ground Corn, Rice Bran, Calcium Carbonate, Wheat Germ, Linseed Meal, Sucrose, Dried Whey Solubles, Cane Molasses, Salt, Dicalcium Phosphate, Monocalcium Phosphate, Colored with Iron Oxide, Zinc Sulfate, Soybean Oil, Vitamin A Supplement, Calcium Pantothenate, Vitamin D3 Supplement, Natural Flavor, Artificial Flavors, Vitamin B12 Supplement, Riboflavin Supplement, Vitamin E Supplement, Copper Sulfate, Niacin Supplement, Manganese Sulfate, Zinc Oxide, Manganous Oxide, Ferrous Sulfate, Potassium Iodide, Sodium Molybdate, Cobalt Carbonate, Sodium Selenite.

DIRECTIONS:

HORSES -

FOALS: Supplement the starting creep ration with 10% to 12% ANIMAX® (200 lbs to 250 lbs per ton) starting on third day and continuing until foal is a year old.

YEARLINGS AND MATURE HORSES: 0.5 lb per horse per day replacing an equal amount of the regular ration.

BROOD MARES: 1 lb to 1.5 lbs ANIMAX[®] daily, starting 3 months before foaling and continuing throughout lactation.

STALLIONS: 1 lb to 1.5 lbs daily with the regular ration.

CATTLE -

DAIRY CALVES: Mix 1 lb ANIMAX® with 1.5 to 2 lbs of cracked or rolled corn. Feed free choice along with water, but small amounts daily to keep feed fresh. Wean at 4 to 5 weeks at which time calves should be eating 1.5 to 2 lbs daily. Continue to feed free choice along with water until 3 months of age. Then feed good quality hay and water free choice along with 4 to 6 lbs of this mix until 7 months of age.

MILK COWS: Contact your Purina® representative for feeding program.

BEEF COWS: to creep feed beef calves, formulate a ration containing 10% to 15% ANIMAX® (200 to 300 lbs per ton) with grain or regular ration and feed until calves are weaned (or longer for extra bloom and finish). For show animals, feed 1 lb to 2 lbs of ANIMAX® per head daily.

SWINE -

SHOW HOGS: 1. Feed 0.25 to 0.5 lb of ANIMAX® with daily meal of show chow for extra show condition. 2. Feed 0.5 to 1 lb of ANIMAX® per day, replacing an equal amount of the regular show chow to hold show condition.

BROOD SOWS: Feed 0.25 lb daily starting two weeks before farrowing.

LACTATING SOWS: Top-dress 0.5 to 1 lb daily in first 3 to 4 days after farrowing.

OTHER ANIMALS -

SHOW TURKEYS: Feed 1 to 2 tablespoons/bird/day or 2 to 4 lbs/100 birds. Blend with pellets.

SHOW CHICKENS: Feed 1 tablespoon/bird/day or 2 lbs/100 birds. Blend with pellets.

SHOW GAME BIRDS: Feed 1 tablespoon/bird/day or 2 lbs/100 birds. Blend with pellets.

OTHER ANIMALS: For directions on feeding ANIMAX® to rabbits or goats, contact your local Purina® dealer.

Available Options:

Product No.	Package	Product Form	Feeding Rate
0001179	Bag 50#	5/32" pellet	See feeding directions above

Product Features:	Product Benefits:
28% protein supplement	Economical, highly concentrated form
Formulated for many different livestock species	No need to separate supplements
Contains milk products and other highly nutritious ingredients	Formulated with nutrients to meet needs of animals in easily digested form
Concentrated source of essential proteins, vitamins, minerals and high energy ingredients	Supports optimal growth and efficient gains

IMPORTANT:

Provide fresh, clean water near the feeding area.

CAUTION:

Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death.