

## BLUE RIBBON LAMB & KID ELECTROLYTE PACK™

with SPECIAL DIRECT-FED MICROBIALS\*



### DESIGNED FOR LAMBS & KIDS

BLUE RIBBON LAMB & KID ELECTROLYTE PACK is specifically designed for lambs and kids as a supplemental source of electrolytes/nutrients.

### A WINNING COMBINATION

BLUE RIBBON LAMB & KID ELECTROLYTES are formulated to provide a supplemental source of electrolytes/nutrients and direct-fed microbials in one product.

### FEATURES

- Highly palatable.
- Contains specifically balanced electrolytes.
- Contains glucose and glycine.
- Contains sodium bicarbonate.
- Contains specifically designed direct-fed microbials at guaranteed levels.

### GUARANTEED ANALYSIS

Sodium, minimum	7.90%
Potassium, minimum	0.45%
Calcium, minimum	0.15%
Calcium, maximum	0.35%
Magnesium, minimum	0.08%
Total Lactic Acid Bacteria	582,010 CFU per g

\*a source of live, viable, naturally occurring microorganisms.

### INGREDIENTS

Glucose, sodium bicarbonate, glycine, potassium chloride, sodium chloride, calcium lactate, magnesium sulfate, citric acid, ascorbic acid, sodium silico aluminate, [dried Lactobacillus acidophilus fermentation product, dried Lactobacillus lactis fermentation product, dried Lactobacillus casei fermentation product, dried Streptococcus diacetylactis fermentation product, dried Bifidobacterium bifidum fermentation product], and artificial coloring (FD & C Blue No. 1).

### PACKAGING

Available in 3 lb pails.

### PRODUCT NO.

32940507 Case: 4-3 lb pails

### Recommended Dose Size By Body Weight

<u>Under 8 lbs</u>	<u>8-12 lbs</u>	<u>12-20 lbs</u>	<u>Over 20 lbs</u>
4-6 oz	8 oz	12 oz	16 oz

### MIX AND FEED INSTRUCTIONS

Mix one measure (rounded scoop - 10 grams) of BLUE RIBBON with each 8 ounces of 90°-100° F water. Feed BLUE RIBBON electrolytes 2-4 hours after milk/milk replacer feeding. BLUE RIBBON electrolytes should not be added to milk/milk replacer. Discard any solution not consumed in 12 hours. Follow veterinary advice for treatment of disease factors. If you have any questions, consult Merrick's Technical Services at 1-800-458-0194 or 1-800-MER-RICK (637-7425).

	DAY 1	DAY 2	DAY 3	DAY 4
For lambs/kids that have been moved or transported:	1 dose BLUE RIBBON 2X daily + 1 feeding MERRICK's milk replacer 2X daily	1 dose BLUE RIBBON 2X daily + 1 feeding MERRICK's milk replacer 2X daily	MERRICK'S milk replacer	MERRICK'S milk replacer
For sick lambs/kids:	1 dose BLUE RIBBON 3-4X daily 4th feeding recommended + 1 feeding MERRICK's milk replacer 2X daily	1 dose BLUE RIBBON 2-3X daily 3rd feeding recommended + 1 feeding MERRICK'S milk replacer 2X daily	1 dose BLUE RIBBON 1-2X daily 2nd feeding recommended + 1 feeding MERRICK'S milk replacer 2X daily	MERRICK'S milk replacer
For weaned lambs/kids:	1 dose BLUE RIBBON 3-4X daily	1 dose BLUE RIBBON 2-3X daily	1 dose BLUE RIBBON 1-2X daily	1 dose BLUE RIBBON 1X daily