

**HERBICIDE****WILBUR-ELLIS**

FOUNDATION[®]

TURF HERBICIDE

2,4-D DICAMBA TRICLOPYR	GROUP	4	HERBICIDE
SULFENTRAZONE	GROUP	14	HERBICIDE

USE THIS PRODUCT ONLY IN ACCORDANCE WITH ITS LABELING AND
WITH THE WORKER PROTECTION STANDARD, 40 CFR PART 170.

ACTIVE INGREDIENTS:

Triclopyr, triethylamine salt	7.28%
Sulfentrazone	0.63%
2,4-D, diethanolamine salt	30.84%
Dicamba, dimethylamine salt	2.48%

OTHER INGREDIENTS:	58.77%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 0.50 lb. 3,5,6-trichloro-2-pyridinyloxyacetic acid per gallon or 5.22%.
 - 0.06 lb. N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.63%.
 - 2.00 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 20.90%.
 - 0.20 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 2.06%.
- Isomer Specific by AOAC Methods.

*Not for sale, distribution or
use in Nassau or Suffolk
Counties in New York State.*

EPA Reg. No. 2217-921-2935

EPA Est. No. 2217-KS-1

KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que
se la explique a usted en detalle.

(If you do not understand the label, find someone to
explain it to you in detail.)

See inside pages for complete Precautionary Statements and
Directions for Use, including Agricultural Use Requirements
and Storage and Disposal.



2023-0718 • 843/5-2023 • AP042423

NET CONTENTS: 2½ GALLONS (9.46 Liters)



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Human and Domestic Animals

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear such as safety glasses. Harmful if swallowed. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

All mixers, loaders, applicators and other handlers must wear:

- protective eyewear,
- long-sleeved shirt and long pants,
- shoes and socks, plus
- waterproof gloves (except for applicators using ground boom equipment) and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical information.	

Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to plants, and to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Groundwater Advisory: These chemicals (triclopyr, 2,4-D and dicamba) have properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Application around a cistern or well may result in contamination of drinking water or groundwater.

Non-target Organism Advisory Statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by minimizing spray drift.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several weeks after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of triclopyr from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms (including sod farms), forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls,
- chemical-resistant gloves made of any water-proof material,
- shoes plus socks and
- protective eyewear.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Reentry Statement: Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

1. Product Description

Designed for turfgrass applications, non-crop, and IVM (Industrial Vegetation Management) applications, this product contains four active ingredients.

- (1) Triclopyr provides broad-spectrum weed control for some of the tough broadleaf weeds such as wild violet, ground ivy, oxalis and wild blackberry.
- (2) Sulfentrazone causes desiccation and yellowing of plant tissue on emerged, susceptible weeds. Sulfentrazone is in the aryl triazolinone family and inhibits protoporphyrinogen oxidase (Protox), a pivotal enzyme in chlorophyll production. Without this key enzyme, a build-up of peroxide-like compounds occurs, thus causing the plant cell membranes of weeds to rupture. Sulfentrazone provides post emergent weed control for common weed species in turfgrass such as spurge and thistles and suppression of yellow nutsedge.

- (3) 2,4-D is an auxin-type herbicide, which is a class of plant growth regulators. It is absorbed through the leaves and is translocated to the growing points of the plant, causing weed stems curl and twist, leaf cupping and withering, and eventual plant death.
- (4) Dicamba is absorbed through the leaves and roots and has multiples modes of actions for hard-to-kill broadleaf weeds.

Combining these herbicides provides a very wide spectrum of weed control for tough and susceptible weeds. FOUNDATION® TURF HERBICIDE controls weeds by affecting multiple sites within the broadleaf weeds. The symptoms of susceptible broadleaf weeds include leaf and stem curl or twisting, and weed yellowing.

These combined herbicides provide limited residual activity.

FOUNDATION TURF HERBICIDE offers these advantages:

- Excellent postemergent activity with proven performance.
- This product exhibits improved cool-weather performance compared to standard “3-way amines”.
- High selectivity (turfgrass safety) in established cool-season turfgrass.
- Sulfentrazone combinations provide rapid and effective weed control for common and troublesome (tough) weed species in turfgrass, including: dandelion, spurge and white clover.
- The speed of action (rate of weed phytotoxicity) and the early weed symptoms are features of sulfentrazone combinations compared to standard “3-way amines”. Often, the weed injury symptoms can be noticed within hours of the application and plant death can occur within 10 to 14 days.
- Triclopyr combinations broaden the weed control spectrum to include many woody and hard-to-control species.
- This product is generally rainfast in as little as 6 hours.

2. Use Restrictions

- The maximum single application rate for FOUNDATION TURF HERBICIDE to ornamental turfgrass, sod farms and non-cropland is 4 pints of product per acre per application, the equivalent of, 1.00 lb 2,4-D ae, 0.25 lb triclopyr ae 0.10 lb dicamba ae, and 0.03 lb sulfentrazone ai per acre per application.
- The maximum annual application rate for FOUNDATION TURF HERBICIDE to ornamental turfgrass and sod farms is 8 pints of product per acre per year, the equivalent of, 2.00 lb 2,4-D ae, 0.50 lb triclopyr ae 0.20 lb dicamba ae, and 0.06 lb sulfentrazone ai per acre per year.
- The maximum number of broadcast applications for commercial sod farms is limited to 2 per year with a minimum of 28 days between applications.
- The maximum number of broadcast applications for ornamental turfgrass is limited to 2 per year.
- The maximum number of broadcast applications to woody plants in non-cropland areas is limited to 1 per year.
- The maximum number of broadcast applications to annual & perennial weeds in non-cropland is limited to 2 per year with a minimum of 30 days between applications.
- Do not apply this product to St. Augustinegrass, bentgrass, greens or tees, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not broadcast apply this product when temperatures are above 90°F, some injury may be expected with spot treatments when air temperatures exceed 90°F. Where state, county or local governments have more stringent temperature regulations, these regulations must be observed.
- To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, nematodes, improper mowing or

improper applications of fertilizer and pesticides. Injury can occur if this product is applied under any of these or other stress conditions. Under any of these stress conditions, any turf damage caused by the use of this product is beyond the control of the registrant and all risk is assumed by the buyer and/or user.

- Do not apply product to bare ground.
- For ground application only; aerial applications are not permitted.
- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (non-cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays).
- Do not apply to wetlands (swamps, bogs, potholes, or marshes).
- Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks and canals.
- Do not apply to agricultural drainage water or on agricultural ditchbanks.
- Do not apply to sites being used for commercial timber or seed production, or to other non-turf plants being grown for sale or other commercial use.
- This product is persistent and may be present in treated plant materials for over 30 days after application. Do not sell or transport treated plant materials or manure from animals that have grazed on treated plant materials off-site for compost distribution or for use as animal bedding/feed for 30 days after application.
- Do not apply or allow this product to come into direct contact with cotton, grapes, tobacco, vegetable crops, flowers, fruit or ornamental trees, or other desirable broadleaf plants; small amounts of spray drift may injure susceptible plants, including ornamental trees or shrubs.

3. Weed Resistance Management

For resistance management, this product contains Group 4 and Group 14 herbicides. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 or Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 4 or Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout area prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with

controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.
- For further information or to report suspected resistance, call 877-800-5556.

4. Spray Preparation And Tank Mixtures

In certain applications, liquid fertilizer may replace part of the water.

Mixing with water:

Add one-half the required amount of water to the spray tank, then add FOUNDATION TURF HERBICIDE slowly with agitation, and complete filling the tank with water. Mix thoroughly and continue agitation while spraying.

When this product is left standing for extended periods of time, re-agitate to assure uniformity of the spray mixture.

Do not use tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer the spray solution to alter the pH range as appropriate.

Mixing with liquid fertilizers:

Use suitable sources and rates of fertilizer based upon recommendations of your fertilizer supplier or State Extension Service Specialist.

Verify physical compatibility with a jar test: Always perform a jar test for compatibility before large scale mixing. The jar test can be conducted by mixing all components in a small container in proportionate quantities. If the mixture separates after standing and can be mixed readily by shaking, then the mixture can be used and applied with spray equipment providing continuous agitation. If large flakes, sludge, gels or other precipitates form, or if a separate oily layer or oil globules appear, then the herbicide and the liquid fertilizer must not be prepared as a tank mixture.

Liquid fertilizers are either solutions (true fluids) or suspensions. Mixing this product with suspensions or N-P-K solutions may not be satisfactory (may be marginal) without pre-mixing this product with water. Premixing this product with 2 parts water will ensure that the emulsifiers are activated enabling the herbicide to be suspended in the fertilizer.

Adjuvants and spray additives:

Adjuvants (such as surfactants, spreaders, spreader-stickers, spray thickeners, foaming agents, activators, detergents, and drift reducing agents) combined with this product can damage the leaf tissue of turfgrass. If any discoloration or cosmetic effects are objectionable or would be unacceptable, then adjuvant(s) combined with FOUNDATION TURF HERBICIDE would not be recommended. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury. When an adjuvant is used with this product, Wilbur Ellis recommends the use of an adjuvant certified by the Council of Producers and Distributors of Agrotechnology (CPDA).

5. Ground Equipment

Spray distribution: The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator. Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. Boom sprayers equipped with appropriate nozzles, tips, and screens are suitable for broadcast applications. For best spray distribution and coverage, select a spray volume and delivery system that will ensure accurate and uniform coverage.

Spray volumes of 10 to 220 gallons per acre with spray pressures adjusted to between 20 to 40 psi. Use higher spray volumes for dense weed populations (up to 220 gallons per acre or 5 gallons per 1,000 square feet).

- Calibration and proper application are essential when using this product.
- Over-application or rates above those specified on this label can cause turf injury.
- Hand-held technique: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.

Hand operated sprayers including backpack sprayers, and compression sprayers are appropriate for small turfgrass areas.

After using this product, clean sprayer with soap or detergent and water, or an approved spray tank cleaner and rinse thoroughly before applying other pesticides.

6. Where To Use

This product provides broadleaf control in the following sites.

- **Ornamental Turfgrass sites:**
 - **Residential/domestic sites** are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
 - **Ornamental turf sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
 - **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs), and office buildings.
- **Non-cropland sites:** including farmyards, fencerows or fence lines, highway rights-of-way (principal, interstate, county, private, and unpaved roads): Roadsides, roadside ditches, road shoulders, road embankments, dividers, and medians; Industrial sites: Lumberyards, tank farms, fuel or equipment storage areas; Municipal, state, and federal lands: Airports and military installations; railroad rights-of-way, railroad yards, railroad crossings and railroad bridge abutments; Utility rights-of-way: telephone, pipeline, electrical powerlines, and communication transmission lines.
- **Agricultural site:** Commercial sod production

Turfgrass tolerance:

- The turfgrass tolerance to this product may vary and temporary turfgrass yellowing may occur. Adverse environmental conditions may reduce the selectivity on the turfgrass. Do not apply this product to stressed turf.

- Certain spray tank additives (adjuvants, wetting agents, surfactants), liquid fertilizers, and tank mixtures containing emulsifiable concentrates may reduce the selectivity on the turfgrass. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury. When an adjuvant is used with this product, Wilbur Ellis recommends the use of an adjuvant certified by the Council of Producers and Distributors of Agrotechnology (CPDA).

STATE RESTRICTIONS:

- **California:** Make broadcast applications only between March 1 and September 1. If troublesome weeds appear during other times of the year, a spot application can be made. While irrigation is necessary and important for plant growth, apply irrigation water efficiently so that no more than 125% of the net irrigation requirement is applied for any irrigation event. Apply efficient irrigations for six months following application of sulfentrazone containing products.
- **Arizona:** The state of Arizona has not approved this product for use on sod farms.
- **New York:** Only one application per year of this product is allowed. This product is not allowed to be sold, distributed or used in Nassau or Suffolk Counties.

7. Mandatory Spray Drift Management

Ground Boom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to select nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Boomless Ground Applications

- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Select medium to very coarse droplet size when sulfentrazone is used postemergence with a contact burndown herbicide. Do not apply as spray droplets smaller than medium to coarse (defined by the ASABE* standard).

*ASABE – American Society for Agricultural and Biological Engineers.

Controlling Spray Droplet Size – Ground Boom

Volume: Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.

Pressure: Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.

Spray Nozzle: Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Boom Height - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

Shielded Sprayers: Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with uniform deposition of the spray on the target area.

Temperature and Humidity: When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions: Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Boomless Ground Applications: Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications: Take precautions to minimize spray drift.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Additional requirements for ground boom application: Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Spray Volume

Ground applicators must use a minimum finished spray volume of 10 gallons per acre. When sulfentrazone is tank mixed with a contact burndown herbicide, ground applicators must use a minimum spray volume of 15 gallons per acre.

8. Application Schedules

Apply this product to broadleaf weeds that are young and actively growing for the best results. Fall applications can provide improved control for emerged winter annuals and perennials such as henbit, chickweed, clover and ground ivy.

For the Listed Residential/domestic sites, Ornamental Turf sites, Institutional sites and Agricultural sites:

Do not apply more than 2 broadcast treatments of this product per site per year. A second broadcast application or a follow-up application as a spot treatment is recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are actively growing.

For the Listed Non-cropland sites:

Use only one (a single) broadcast treatment of this product per site per year for woody plant infestations, or two broadcast treatments for annual and perennial weeds. It is required to wait 30 days between treatments. A spot treatment may be substituted for the second broadcast treatment if necessary.

Extremes in environmental conditions e.g. temperature and moisture, soil conditions, and cultural practices may affect the activity of this product. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms is delayed, and weeds hardened off by drought are less susceptible to this product.

For newly seeded areas:

Delay application of this product to grass seedlings until after the second mowing.

For newly sodded, sprigged, or plugged areas:

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Reseeding interval:

Treated areas may be reseeded 3 weeks after application.

9. How Much To Use

USE RATES AND SPRAY VOLUMES:

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

TABLE 1. USE RATES FOR ORNAMENTAL TURF SITES, NON-CROPLAND AND SOD FARMS		
Species	Rate	Recommended Spray Volume
Cool-season Turf		
Kentucky bluegrass, annual bluegrass, annual ryegrass perennial ryegrass, tall fescue, red or fine leaf fescues	3.25 to 4 Pints/Acre (1.2 to 1.5 fl.oz./1,000 sq.ft.)	10 to 220 Gallons/Acre (0.25 to 5.0 Gallons/1,000 sq.ft.)

**SPOT TREATMENT:
WITH HAND OPERATED SPRAYERS (INCLUDING BACKPACK SPRAYERS AND
PUMP-UP TYPE SPRAYERS):**

- Apply any time the emerged broadleaf weeds are actively growing.
- Calibration and proper application are essential when using this product.
- Uniform applications are essential when using this product. Over application or rates above those specified on this label including excessive overlaps of this product can cause turf injury.
- Hand-held techniques: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.
- Follow-up applications as spot treatments are recommended for more mature weeds, for dense infestations, and for adverse environmental conditions. Wait 30 days before making a follow-up application.
- **For cool-season turfgrass listed in Table 1:** Mix 1.2 to 1.5 fl.oz. of this product per 1 gallon of water for treatment of approximately 1,000 sq.ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.

10. Cultural Tips

Irrigation:

- Do not apply this product through any type of irrigation system.
- Rainfast in as little as 6 hours. Do not apply this product immediately before rainfall or irrigation.
- If possible, do not irrigate or water the turfgrass within 6 to 24 hours after application.
- If dry conditions exist, a scheduled irrigation or watering 24 hours before and 24 hours after application is recommended.

Mowing:

- For optimum results, delay mowing 2 days before and until 2 days after the application of this product.

11. Broadleaf Weeds Controlled

FOUNDATION TURF HERBICIDE will control or suppress the following broadleaf weeds. Apply any time the emerged broadleaf weeds are susceptible.

BROADLEAF WEEDS			
Aster, white heath & white prairie	False dandelion (*spotted catsear & common catsear)	Mallow, common	Redweed
Bedstraw	Field bindweed	Matchweed	Red sorrel (*sheep sorrel)
Beggarweed, creeping	(*morningglory & creeping jenny)	Mouseear chickweed	Russian Thistle
Bindweed	Field oxeye-daisy (*creeping oxeye)	Mustard	Shepherd's purse
Black medic	Filaree, whitestem & redstem	Nettle	Speedwell (Veronica)
Broadleaf plantain	Florida betony	Nutsedge** (yellow)	Spurge
Buckhorn plantain	Florida pusley	Old world diamond flower	Thistle
Bull thistle	Ground ivy	Oxalis (*yellow woodsorrel, creeping woodsorrel, and corniculata)	Virginia buttonweed
Burdock, common	Groundsel	Parsley-piert	White clover (*Dutch clover, honeysuckle clover, white trefoil, & purplewort)
Buttercup, creeping	Hawkweed	Pennsylvania smartweed	Wild carrot
Carpetweed	Healall	Pepperweed	Wild garlic
Catnip	Henbit	Pigweed	Wild geranium
Chickweed	Innocence (Blue-eyed Mary)	Pineappleweed	Wild lettuce
Chicory	Knotweed	Plantain	Wild mustard
Cinquefoil	Kochia	Poison ivy	Wild onion
Clover	Lambsquarters	Poison oak	Wild strawberry
Curly dock	Lawn burweed	Prickly lettuce (*compass plant)	Wild violet***
Dandelion	Lespedeza, common	Puncturevine	Yarrow
Dayflower		Purple cudweed	Yellow rocket
Deadnettle		Purslane	
Dock		Ragweed	
Dogfennel			
English Daisy			
<p>* Synonyms ** Suppression only when nutsedge is young and actively growing. *** For best results, apply in the spring when wild violets are blooming or apply a late fall application followed by a spring application.</p>			

12. For Use In Non-Cropland: Brush And Weed Control

High volume foliar applications (100 to 400 gallons per acre):

Apply 0.5 to 2.0 gallons of product per acre with adequate water or apply a 0.5 to 2.0% vol/vol spray solution as a full cover spray with high volume equipment. Use the lower spray concentrations in the range for susceptible species and use the higher spray concentrations within the range for hard-to-control species, for mature plants during the late summer or under adverse environmental conditions (e.g. drought).

Spray broadleaf weeds, woody plants or mixed brush uniformly and thoroughly by wetting all leaves, stems, bark and root collars. The total volume of spray solution required for adequate coverage of solid stands of mixed brush can range from 100 to 400 gallons of spray solution per treated acre. The spray preparation chart for applications on a spray-to-wet basis is shown below in Table 2.

Table 2. Instructions for preparing 100 to 400 gallons of spray solution at 0.5 to 2.0% spray concentration with water for high volume foliar applications.

Spray solution per acre, Gallons	Amount of Product Needed for Spray Concentration of:			
	0.5%	0.67%	1.0%	2.0%
100	0.5 gal	0.67 gal	1.0 gal	2.0 gal
200	1.0 gal	1.34 gal	2.0 gal	—
300	1.5 gal	2.0 gal	—	—
400	2.0 gal	—	—	—

Equal measures: 1 gallon = 4 quarts = 8 pints = 128 fl.oz.

The maximum seasonal application rate for trees, brush and woody plant control is 2.0 gallons of product per acre per application per site.

For Backpack Sprayers and Hand Pump-up Sprayers.

Table 3. Instructions for preparing 1 to 3 gallons of spray solution at 0.5 to 2.0% spray concentration with water for high volume foliar applications.

Gallons of Water	Amount of Product Needed for Spray Concentration of:			
	0.5%	0.67%	1.0%	2.0%
1	4 teaspoons	5 teaspoons	2.5 tablespoons	5 tablespoons
2	2.5 tablespoons	3.5 tablespoons	5 tablespoons	10 tablespoons
3	2 fl.oz.	2.5 fl.oz.	4 fl.oz.	8 fl.oz.

Equal measures: 1 fl.oz. = 2 tablespoons (Tbs.) = 6 teaspoons (tsp.)

BRUSH CONTROLLED

Ash	Cedar	Honeylocust (suppression)	Sycamore
Aspen	Cherry (except Black Cherry)	Honeysuckle	Trumpet creeper
Beech	Cottonwood	Kudzu	Wild Grape
Birch	Dogwood	Multiflora Rose	Wild Plum
Blackberry	Elm	Oak	Willow
Black Locust	Gooseberry	Poison Ivy	
Brambles	Hawthorn	Poison Oak	
Buckbrush		Sumac	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons:

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For Plastic Containers – Nonrefillable with capacities greater than 5 gallons:

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(cont. on next page)

STORAGE AND DISPOSAL *(cont.)*

For Refillable Containers:

CONTAINER HANDLING: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Container cleaning: Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.


ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

 NOTICE: This product can expose you to diethanolamine, which is known to the State of California to cause cancer. For more information go to www.P65Warnings.ca.gov.

WILBUR-ELLIS® logo, FOUNDATION® and HERBICIDE® logo are registered trademarks of Wilbur-Ellis Company LLC.

843/5-2023

2023-0718 • AP042423

**For Chemical Spill, Leak, Fire or Exposure,
Call CHEMTREC (800) 424-9300**

Manufactured for:

Wilbur-Ellis Company LLC

2903 S. Cedar Avenue • Fresno, CA 93725

THIS PAGE IS BLANK INTENTIONALLY.

THIS PAGE IS BLANK INTENTIONALLY.



HERBICIDE



WILBUR-ELLIS

FOUNDATION® TURF HERBICIDE

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

ACTIVE INGREDIENTS:

Triclopyr, triethylamine salt	7.28%
Sulfentrazone	0.63%
2,4-D, diethanolamine salt	30.84%
Dicamba, dimethylamine salt	2.48%

OTHER INGREDIENTS: 58.77%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

- 0.50 lb. 3,5,6-trichloro-2-pyridinyloxyacetic acid per gallon or 5.22%.
- 0.06 lb. N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl] methanesulfonamide per gallon or 0.63%.
- 2.00 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon on 20.90%.
- 0.20 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 2.06%.

Isomer Specific by AOAC Methods.

EPA Reg. No. 2217-921-2935

EPA Est. No. 2217-KS-1

KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Human and Domestic Animals

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear such as safety glasses. Harmful if swallowed. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

All mixers, loaders, applicators and other handlers must wear: protective eyewear, long-sleeved shirt and long pants, shoes and socks, plus water-proof gloves (except for applicators using ground boom equipment) and chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate. See engineering controls for additional requirements.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

2,4-D DICAMBA TRICLOPYR	GROUP	4	HERBICIDE
SULFENTRAZONE	GROUP	14	HERBICIDE

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical information.

843/5-2023

2023-0718 • AP042423

Manufactured for: **Wilbur-Ellis Company LLC**
2903 S. Cedar Avenue • Fresno, CA 93725

NET CONTENTS: 2½ GALLONS (9.46 Liters)